

# **Astrophysics to Understand the Universe Mass Density, Vacuum Energy Density, and Curvature**







#### **Fundamental Questions:**


- Will the universe last forever?
- *Is the universe infinite?*
- What is the universe made of?

#### An unusual moment in human history:

At the beginning of this century, Einstein developed the conceptual tools to address these questions empirically.

In the past decade or so, technology has advanced to the point that we can now make the measurements that begin to answer these fundamental questions.

Progress is now being made with large scientific programs, including the Supernova Cosmology Project and the Cosmic Microwave Background satellites: MAP, and PLANCK.



#### The implications of an accelerating universe:


- 1. The expansion is not slowing to a halt and then collapsing (i.e., the universe is *not* "coming to an end"). In the simplest models, it will expand forever.
- 2. There is a previously unseen energy pervading all of space that accelerates the universe's expansion.

This new accelerating energy ("dark energy") has a larger energy density than the mass density of the universe (or else the universe's expansion wouldn't be accelerating).

#### What we don't know is:

- 1. How much of mass density and dark energy density is there? I.e., how much dark matter and dark energy do we need to look for?
  - The answer to this question determines the "curvature" of the universe, and can tell us about the extent of the universe: infinite or finite.
- 2. What is the "dark energy"? Particle physics theory proposes a number of alternatives, each with different properties that we can measure. Each of the alternative theories raises some important questions/problems of fundamental physics.









## supernova / acceleration probe satellite overview

- 1.8m aperture telescope

  Can reach very distant SNe.
- 1 square degree mosaic camera, 1 billion pixels

  Efficiently studies large numbers of SNe.
- 3-channel spectroscopy, 0.3um -- 1.8um Detailed analysis of each SN.

#### MIDEX+ class satellite:

Dedicated instrument.

Designed to repeatedly observe an area of sky.

Essentially no moving parts.

4-year construction cycle.3-year operation for experiment (lifetime open-ended).



## supernova acceleration probe satellite baseline specifications

Aperture 1.8	3m
--------------	----

F.O.V. 1 square degree
Launch Vehicle Delta II 7920

Image array	36k x 36k mosaic CCD
Architecture	2k x 4k 10-micron pixels
Plate Scale	10 micron/0.10 arc-sec

Image	Arrass	Banda	D 17 D T 7
ııııaqe	Allay	Banas	B,V,R,I,Z

**Q.E.** 65% @ 1000nm (LBNL CCD)

92% @ 900nm

>85% @ 400nm-800nm

Read Noise <2-4 e-

Image Array Temp. 150K

Spectroscopy	3-channel	<pre>pick-off:</pre>
--------------	-----------	----------------------

325-600nm "Blue" CCD

+ 600-1050nm "Red" LBL CCD

+ 1050-1800nm IR (HgCdTe)

Thermal Cooling passive

Data Downlink 320 Mbps

Data Storage 256 Gbits solid state

Pointing Stability <0.1 arc-sec
Image Stabilization 2-axis fast</pre>

steering mirror

Satellite Mass <1500kg
Power Consumption <800W



#### supernova acceleration probe one-year baseline data package

#### Full sample of 2000 SNe between z = 0.3 and 1.7

Discovery within ~2 days of explosion (i.e. ~2 weeks before max). Most dense coverage between z = 0.3 and 1.0

- Spectra at max for all SNe. (0.3 -- 1.8um)
- Lightcurve points at least 1/week (restframe) from -15 to +60 days (restframe)

#### Discover every SN in the field


Nearly continuous monitoring of ~2 sq. deg. to  $m_{AB}$  (@1um)  $\approx$  28.5+ ~10 sq. deg. to  $m_{AB}$  (@1um)  $\approx$  25+ ~100 sq. deg. to  $m_{AR}$  (@1um) ≈ 24+

#### Satellite follows additional 200 SNe at z < 0.15

Discovery from ground, follow-up from space to ensure matching spectroscopy/photometry across redshifts.

#### Subsample of 200 SNe from the full sample

- Selected to span lightcurve timescales
  - galaxy environments (morphology, galactocentric radius)
  - redshifts
- Spectra at least 2/week (restframe) first month 1/week (restframe) later
- Synthetic "filter-tuned" photometry from spectra for perfect K-corrections





Target

**Target** 

Parameter

Constraint

Statistical Uncertainty

#### Determine mass density, vacuum energy density, and curvature

mass/vacuum $\Omega_M$ ,	•	in flat universe $\Omega_k$ =0		0.01	
curvature	$\Omega_k$	(indep. of CMB	)	0.05	
mass density vacuum energy	$\Omega_M$ y $\Omega_{\Lambda}$	unconstrained	$\Omega_k$	0.02 0.05	

#### **Properties of Dark Energy**

Eq. of State w vs  $\Omega_M$  in flat universe


Eq. of State w with  $\Omega_M \approx 0.3$ 

0.05


Study time-varying w(t) by studying  $d_L(z)$  with  $\Delta z \approx 0.03$  bins.

A definitive supernova cosmology measurement.


Supernova Cosmology Project Perlmutter *et al.* (1998)



Supernova Cosmology Project Perlmutter *et al.* (1998)




# Dark Energy Unknown Component, $\Omega_{u}$ , of Energy Density



SnapSat
Target Statistical Uncertainty

# Dark Energy Unknown Component, $\Omega_{u}$ , of Energy Density



SNAP Satellite
Target Statistical Uncertainty



#### **Dust and Extinction**

Determine individual extinctions laws for a given supernova, using range of color measurements into near IR.

Determine gray dust contribution (if any), tracking  $d_I(z)$  to high z.

#### **SN Progenitor Age and Metallicity Effects**

Studies of lightcurve-timescale (and spectral) luminosity indicators with significant sample of SNe in all host galaxy environments.

#### **Measurement Systematics**

Observe all SNe at all redshifts with single, calibrated, stabile photometry/spectroscopy system.

(Avoid multiple instruments with different atmospheric/moon conditions.)



## supernova acceleration probe complementary science

#### **Cosmological Parameters...**

Type II supernova expanding photosphere,  $\Omega_M$ ,  $\Omega_\Lambda$  at  $z\approx 0.5$  Strong lensing statistics.  $\Omega_\Lambda$  Weak lensing Galaxy clustering, P(k)

z > 1 clusters and associated lensing

. . .

#### ...and Beyond

GRB optical counterparts: rates, lightcurves, and spectra MACHO optical counterparts by proper motion Galaxy populations and morphology to co-added  $m \approx 32$  Target selection for NGST Kuiper belt objects Supernova rates, star formation rates Supernova phenomenology studies Low surface brightness galaxies, luminosity function

. . .

### Roadmap for the Supernova Cosmology Project


	Vearby SM Factory	Subatu Key Proj	set Nazi	roject SNAPSat
SNe / Year				
@ $z < 0.06$	100	•	•	100
@ $z \sim 0.5$	•	12	•	1000
@ $z > 1$	•	4	20	900
Duration	2 yrs	3 yrs	3 yrs	2 yrs
Calibrated Candle	<b>/</b>	<b>/</b>	<b>/</b>	<b>/</b>
Reddening Measure	<b>/</b>	<b>/</b>	<b>/</b>	<b>/</b>
Gray Dust Measure	<b>/</b>	<b>/</b>	<b>/</b>	<b>/</b>
Metallicity Measure	•		<b>/</b>	<b>/</b>
Evolution Checks	<b>/</b>			<b>/</b>
Standardized Datasets	S .		•	<b>/</b>


### LBNL CCD Technology

High quantum efficiency from near UV to near IR No thinning, no fringing.

High yield.

Radiation hard.





### NASA Interest in New Astrophysics Probes





"... in the quest to quantify the expansion of the Universe, astronomers may have uncovered a new physical phenomenon -- a kind of vacuum energy they call the **Cosmological Constant. This** exciting discovery may have given astronomers and physicists an unexpected clue about fundamental physics."

Daniel S. Goldin, May 28, 1999 Fermilab Inner Space-Outer Space Conference